

CORN (Field, Corn Grown for Seed, Popcorn, and Sweet Corn)

DISEASE CONTROL		USE RATE	APPLICATION TIMING
- Common Rust - Southern Rust - Anthracnose Leaf Blight - Gray Leaf Spot	- Northern Corn Leaf Blight - Northern Corn Leaf Spot - Southern Corn Leaf Blight - Eye spot	5-7 fl oz/A	For optimum results, begin applications when disease first appears and continue as needed on a 7 – 10 day interval. Use high end of the use rate when disease pressure is high and conditions are favorable for disease development. Apply no later than growth stage R4 (early dough stage). For multiple applications, refer to the guidelines under Resistance Management.
	2(ee) Label - Tar Spot Management		

RESTRICTIONS AND OTHER INFORMATION:

- Do not exceed 7 fl oz of product/A for a single application.
- Do not apply more than 2 applications of product per growing season.
- Do not apply more than 14 fl oz/A of product per acre per year.
- Do not apply more than 0.228 lb ai of flutriafol per acre per year.
- Do not apply more than 2.0 lb ai of azoxystrobin per acre per year.
- An adjuvant may be used with PREEMPTOR EQ Fungicide prior to the V8 corn growth stage and after the VT corn growth stage.
- Restricted Entry Interval (REI): The REI for detasselling field corn and popcorn grown for seed is 5 days. The REI for sweet corn is 3 days. The REI for all other activities is 12 hours.
- Preharvest Interval: Do not apply within 7 days of harvest.




Gray Leaf Spot on Corn

SOYBEANS

DISEASE CONTROL		USE RATE	APPLICATION TIMING
- Aerial Blight - Alternaria Leaf Spot - Anthracnose - Brown Spot - Cercospora Blight and Purple Seed Stain - Frogeye Leaf Spot - Pod and Stem Blight	- Powdery Mildew - Target Spot - Rust Suppression: - White Mold/ Stem Rot - Southern Blight	5-8 fl oz/A	For Cercospora Leaf Blight, begin applications at first flower (R1). For Rust, apply beginning at first visible flower (R1) or when conditions are favorable for disease development. For Aerial Blight, use high rate. For all other diseases, apply preventatively or when conditions are favorable for disease development. Use the higher rate or additional application when environmental conditions are conducive for disease development. For multiple applications, refer to the guidelines under Resistance Management. An adjuvant may be added at recommended rates. Minimum retreatment interval: For soybean rust, the minimum retreatment interval is 21 days; maximum retreatment interval is 35 days. All other diseases: the minimum retreatment interval is 14 days; maximum retreatment interval is 35 days.

RESTRICTIONS AND OTHER INFORMATION:

- Do not feed forage or hay to animals or permit animals to graze.
- Do not apply more than 15.5 fl oz/A of product per acre per year.
- Only one application of product at 8 fl oz/A may be made to any one field during a single growing season.
- Do not apply more than 3 applications per year.
- Do not apply more than 0.23 lb ai flutriafol per acre per year.
- Do not apply more than 1.5 lb ai of azoxystrobin per acre per year.
- Restricted Entry Interval (REI): The REI is 12 hours.
- Preharvest Interval: Do not apply within 21 days of harvest.



Frogeye Leaf Spot on Soybean

Application Timing for Corn: V5-V8/VT-R1



Application Timing for Soybeans: V2-R5



Always consult the product label for complete use directions.